

Discrete Mathematical Structures CS 3233 Lecture 11

Prof. William Winsborough
October 11, 2007

Business

- Homework 7 due Thursday 10/18
 - Section 2.3: 10, 14, 16, 38
 - Section 2.4: 4, 8

11 October 2007

Winsborough CS 3233 Lecture 11

2

Injections and Surjections

- $f : A \rightarrow B$ is *one-to-one*, or *injective*, if and only if $f(x) = f(y)$ implies $x = y$ for all $x, y \in A$
 - $f : A \xrightarrow{1-1} B$
 - $\forall x \forall y. f(x) = f(y) \rightarrow x = y$
 - Equivalently, $\forall x \forall y. x \neq y \rightarrow f(x) \neq f(y)$
 - f is called an *injection*
- $f : A \rightarrow B$ is *onto*, or *surjective*, if and only if for every $b \in B$, there is an $a \in A$ such that $f(a) = b$
 - f is a *function of A onto B*
 - $f : A \xrightarrow{\text{onto}} B$
 - $\forall y \exists x (f(x) = y)$
 - f is called a *surjection*

11 October 2007

Winsborough CS 3233 Lecture 11

3

More Examples

- Injective and/or surjective?
 - Square
 - $f : Z \rightarrow Z$
 - $f(x) = x^2$
 - Double
 - $\text{twice} : Z \rightarrow Z$
 - $\text{twice}(x) = 2x$
 - Absolute value
 - $|\cdot| : Z \rightarrow N$
 - $|x| = \begin{cases} x & \text{if } 0 \leq x; \\ -x, & \text{otherwise} \end{cases}$

11 October 2007

Winsborough CS 3233 Lecture 11

4

Still More Examples

- Injective and/or surjective?
 - Integer successor
 - $zs : Z \rightarrow Z$
 - $zs(n) = n+1$
 - Mapping of 32-bit words to Z
 - Successor in $Z_3 = \{0, 1, 2\}$
 - $s(0) = 1$
 - $s(1) = 2$
 - $s(2) = 0$

11 October 2007

Winsborough CS 3233 Lecture 11

5

Bijections

- If $f : A \rightarrow B$ is injective and surjective, it is *bijective*
 - f is called a *one-to-one correspondence*, or a *bijection*

11 October 2007

Winsborough CS 3233 Lecture 11

6

Compositions

- Definition
 - Given $g : A \rightarrow B$ and $f : B \rightarrow C$, the composition of f and g , $f \circ g : A \rightarrow C$, is defined by $f \circ g(a) = f(g(a))$
 - Note that no special properties of f and g are required for $f \circ g$ to be defined.
 - For instance, f and g need not be injective, surjective, or bijective
 - However, if f and g have special properties, it often follows that $f \circ g$ special properties as well
 - Study hint: think through these relationships

11 October 2007

Winsborough CS 3233 Lecture 11

7

Function Inverses

- Theorem: Given $f : A \xrightarrow{1-1} B$, every $b \in B$ has a unique pre-image $a \in A$
- This justifies the following definition: Given

$$f : A \xrightarrow{1-1} B$$

- $f^{-1} : B \rightarrow A$ is the *inverse* of f
- $f^{-1}(b) = a$ iff $f(a) = b$

11 October 2007

Winsborough CS 3233 Lecture 11

8

Bijections and cardinality

- Sets A and B are *equinumerous* (meaning they have the same cardinality) iff there is a one-to-one correspondence between them
 - Notice that this defines a binary relation over sets
 - Contains the ordered pairs of sets between which there exists a one-to-one correspondence
- Pretty clear for finite sets

11 October 2007

Winsborough CS 3233 Lecture 11

9

Equinumerous Infinite Sets

- \mathbb{N} and \mathbb{Z} have the same cardinality
 - $f : \mathbb{N} \rightarrow \mathbb{Z}$
 - $f(n) = \begin{cases} 0, & \text{if } n = 0 \\ n/2, & \text{if } n \text{ is even} \\ -(n-1)/2, & \text{if } n \text{ is odd} \end{cases}$
 - Note, f is a bijection
 - Use case analysis to show f is injective based on whether x and y are odd or even
- Show \mathbb{N} and \mathbb{Z}^+ are equinumerous

11 October 2007

Winsborough CS 3233 Lecture 11

10

Countability

- Definition
 - A set A is *countable* if it is finite or it is equinumerous to \mathbb{Z}^+
 - Otherwise, A is *uncountable*
- Theorem: \mathbb{Q}^+ , the set of positive rationals, is countable

11 October 2007

Winsborough CS 3233 Lecture 11

11

Uncountability of the Reals

- Theorem
 - \mathbb{R} , the set of real numbers, is uncountable
- Proof
 - Uses Georg Cantor's *diagonalization argument*
 - Outline
 - Assume for contradiction that there is a one-to-one correspondence, f , between \mathbb{N} and the real interval $[0, 1]$
 - Use f to construct a real in $[0, 1]$ that has no preimage under f
 - Idea: for each decimal place, n , in the representation of the constructed value, choose a decimal different from the n^{th} place of $f(n)$
 - The fact that the constructed value differs from each value assumed by f shows that f is not onto, giving the desired contradiction

11 October 2007

Winsborough CS 3233 Lecture 11

12